



RECEIVED

Express Mail No.: EL 500 577 579 US
JUN 23 2003

Sheet 1 of 5

TECH CENTER 1600/2900

| LIST OF REFERENCES CITED BY APPLICANT (Use several sheets if necessary) | | | | ATTY. DOCKET NO. 10624-0092-999 | | APPLICATION NO. 10/009,945 | |
|--|----|---|------------|--|-------|-------------------------------|-------------------------------|
| | | | | APPLICANT Thomsen, G. and Wrana, J. | | | |
| | | | | FILING DATE December 11, 2001 | | GROUP To Be Assigned | |
| U.S. PATENT DOCUMENTS | | | | | | | |
| EXAMINER INITIAL | | DOCUMENT NUMBER | DATE | NAME | CLASS | SUBCLASS | FILING DATE IF APPROPRIATE |
| HA | AA | 6,103,869 | 08/15/2000 | Souchelnnytokyi et al. | 530 | 330 | |
| | AB | 6,087,122 | 07/11/2000 | Hustad et al. | 435 | 29 | |
| | AC | 6,060,262 | 05/09/2000 | Beer-Romero et al. | 435 | 15 | |
| | AD | 6,001,619 | 12/14/1999 | Beach et al. | 435 | 193 | |
| ✓ | AE | 6,194,507 09/30/1998 | 08/30/1999 | Hoekstra et al. | 534 | 441 | |
| FOREIGN PATENT DOCUMENTS | | | | | | | |
| | | DOCUMENT NUMBER | DATE | COUNTRY | CLASS | SUBCLASS | TRANSLATION YES NO |
| | | | | | | | |
| OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.) | | | | | | | |
| | AF | Afrakhte, et al. 1998. "Induction of Inhibitory Smad6 and Smad7 mRNA by TGF- β family members," Biochemical and Biophysical Research Communications 244: 505-511. | | | | | |
| | AG | Baker, et al. 1996. "A novel mesoderm inducer, Madr2, functions in the activin signal transduction pathway," Genes & Development 10:1880-1889. | | | | | |
| | AH | Bartel, et al. 1990. "The recognition component of the N-end rule pathway," EMBO Journal 9: 3179-3189. | | | | | |
| | AI | Bartel, et al. 1995. "Analyzing protein-protein interactions using two-hybrid system," Methods in Enzymology, Vol. 254, (24):1-263. | | | | | |
| | AJ | Bashirullah, et al. 1998. "RNA Localization in Development," Annu. Rev. Biochem. 67:335-94. | | | | | |
| | AK | Bitzer, et al. 2000. "A mechanism of suppression of TGF- β /Smad signaling by NF- κ B/Rel A," Genes & Development 14:187-197. | | | | | |
| | AL | Bonifacino, et al. 1998. "Ubiquitin and the Control of Protein Fate in the Secretory and Endocytic Pathways," Ann. Rev. Cell. Biol. 14:19-57. | | | | | |
| | AM | Chen, et al. 1995. "The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules," Proc. Natl. Acad. Sci. USA 82:7819-7823. | | | | | |
| | AN | Chung, et al. 1998. "A novel, putative MEK kinase controls developmental timing and spatial patterning in Dictyostelium and is regulated by ubiquitin-mediated protein degradation," Genes Dev. 12: 3564-78. | | | | | |
| | AO | Coffman, T.M., 1997. "A genetic approach for studying the physiology of the Type 1A (AT _{1A}) Angiotensin Receptor," Seminars in Nephrology 17:404-411. | | | | | |
| | AP | Derynck, et al., 1998, "Smads: Transcriptional Activators of TGF- β Responses," Cell 19:737-740. | | | | | |
| | AQ | Dickson, B.J., 1998. "Photoreceptor development: Breaking down the barriers," Current Biology 8:R90-R92. | | | | | |
| | AR | Eppert, et al. 1996. "MADR2 Maps to 18q21 and encodes a TGF β -Regulated MAD-Related protein that is functionally mutated in Colorectal Carcinoma," Cell 86: 543-552. | | | | | |

| | |
|----|--|
| AS | Epps, et al. 1998. "The Drosophila semushi mutation blocks nuclear import of Bicoid during embryogenesis," Current Biology 8:1277-1280. |
| AT | Esther, Jr., et al. 1996. "Mice Lacking Angiotensin-Converting Enzyme Have Low Blood Pressure, Renal Pathology, and Reduced Male Fertility," Laboratory Investigation 74:953. |
| AU | Fainsod, et al. 1994. "On the function of BMP-4 in patterning the marginal zone of the Xenopus embryo," Embo J 13:5015-25. |
| AV | Gilboa, et al. 1998. "Oligomeric structure of Type I and Type II transforming growth Factor β Receptors: homodimers form in the ER and persist at the Plasma Membrane," J. Cell Biol. 140:767-777. |
| AW | Govers, et al. 1999. "Identification of a novel ubiquitin conjugation motif, required for ligand-induced internalization of the growth hormone receptor," EMBO J. 18:28-36. |
| AX | Graff, et al. 1996. "Xenopus Mad proteins transduce distinct subsets of signals for the TGF β Superfamily," Cell 86:1-20. |
| AY | Harland, et al. 1997. "Formation and function of Spemann's Organizer," Ann. Rev. Cell Biol. 13:611-667. |
| AZ | Harvey, et al. 1999. "Nedd4-like proteins: an emerging family of ubiquitin-protein ligases implicated in diverse cellular functions." Trends Cell Biol. 9:166-169. |
| BA | Hayashi, et al. 1997. "The MAD-Related protein Smad7 Associates with the TGF β Receptor and Functions as an antagonist of TGF β signaling," Cell 89:1165-1173. |
| BB | Hein, et al. 1995. "NPI1, an essential yeast gene involved in induced degradation of Gap1 and Fur4 permeases, encodes the RspS ubiquitin-protein ligase," Mol. Microbiol. 18:77-87. |
| BC | Heldin, et al. 1997. "TGF- β signalling from cell membrane to nucleus through SMAD proteins," Nature 390:465-71. |
| BD | Hemmati-Brivanlou, et al. 1995. "Ventral mesodermal patterning in Xenopus Embryos: expression patterns and activities of BMP-2 and BMP-4," Dev. Genet. 17:78-89. |
| BE | Hemmati-Brivanlou, et al. 1997. "Vertebrate Embryonic Cells will become nerve cells unless told otherwise," Cell 88:13-17. |
| BF | Henis, et al. 1994. "The Types II and III transforming growth Factor- β Receptors form Homo-Oligomers," J. Cell Biol. 126:139-154. |
| BG | Hershko, et al. 1998. "The Ubiquitin System," Ann. Rev. Biochem. 67:425-479. |
| BH | Hicke, L., 1999. "Gettin' down with ubiquitin: turning off cell-surface receptors, transporters and channels," Trends Cell Biol. 9:107-112. |
| BI | Hochstrasser, M., 1996. "Ubiquitin-Dependent protein degradation," Ann. Rev. Genet. 30:405-439. |
| BJ | Hoodless, et al. 1996. "MADR1, a MAD-Related protein that functions in BMP2 signaling pathways," Cell 85:489-500. |
| BK | Horb, et al. 1997. "A vegetally-localized T-box transcription factor in Xenopus eggs specifies mesoderm and endoderm and is essential for embryonic mesoderm formation," Dev. 124:1689-1698. |
| BL | Huang, et al. 1995. "Control of cell fate by a deubiquitinating enzyme encoded by the fat facets gene," Science 270:1828-31. |
| BM | Huibregtse, et al. 1995. "A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase," Proc. Natl. Acad. Sci. U.S.A. 92:2563-7. |

| | | |
|--|----|--|
| | BN | Imamura, et al. 1997. "Smad6 inhibits signaling by the TGF- β superfamily," <i>Nature</i> 389:622-626. |
| | BO | Ishisaki, et al. 1998. "Smad7 is an activin-inducible inhibitor of activin-induced growth arrest and apoptosis in Mouse B Cells," <i>J. Biol. Chem.</i> 273:24293-24296. |
| | BP | Itoh, et al. 1998. "Transforming growth factor β 1 induces nuclear export of inhibitory Smad7," <i>J. Biol. Chem.</i> 273:29195-29201. |
| | BQ | Jiang, et al. 1998. "Regulation of the Hedgehog and Wingless signalling pathways by the F-box/WD40-repeat protein Slimb," <i>Nature</i> 391:493-6. |
| | BR | Joazeiro, et al. 1999. "The tyrosine kinase negative regulator c-Cbl as a RING-Type, E2-Dependent ubiquitin-protein ligase," <i>Science</i> 286:309-312. |
| | BS | Jonk, et al. 1998. "Identification and functional characterization of a Smad binding element (SBE) in the JunB promoter that acts as a transforming growth Factor- β , activin, and bone morphogenetic protein-inducible enhancer," <i>J. Biol. Chem.</i> 273:21145. |
| | BT | Joseph, et al. 1998. "Mutant Vg1 ligands disrupt endoderm and mesoderm formation in <i>Xenopus</i> embryos," <i>Development</i> 125:2677-85. |
| | BU | Kawabata, et al. 1998. "Signal transduction by bone morphogenetic proteins," <i>Cytokine Growth Factor Rev.</i> 9:49-61. |
| | BV | Kimelman, et al. 1998. "Mesoderm Induction: A Postmodern View," <i>Cell</i> 94:419-21. |
| | BW | Kim, et al. 1997, "Drosophila Mad binds to DNA and directly mediates activation of vestigial by Decapentaplegic," <i>Nature</i> 388:304. |
| | BX | Kretzschmar, et al. 1997. "The TGF- β family mediator Smad1 is phosphorylated directly and activated functionally by the BMP receptor kinase," <i>Genes Dev.</i> 11:984-95. |
| | BY | Kumar, et al. 1997. "cDNA Cloning, expression analysis, and mapping of the Mouse Nedd4 Gene," <i>Genomics</i> 40:435-43. |
| | BZ | Kwon, et al. 1998. "The mouse and human genes encoding the recognition component of the N-end rule pathway," <i>Proc. Natl. Acad. Sci. U.S.A.</i> 95:7898-903. |
| | CA | Levkowitz, et al. 1999. "Ubiquitin ligase activity and tyrosine phosphorylation underlie suppression of growth factor signaling by c-Cbl/Sli-1," <i>Mol. Cell</i> 4:1029-1040. |
| | CB | Lindsay, et al. 1998. "A deubiquitinating enzyme that disassembles free polyubiquitin chains is required for development but not growth in dictyostelium," <i>J. Bio. Chem.</i> 273:24131-8. |
| | CC | Macias-Silva, et al. 1996. "MADR2 is a substrate of the TGF β receptor and its phosphorylation is required for nuclear accumulation and signaling," <i>Cell</i> 87:1215-1224. |
| | CD | Macias-Silva, et al. 1998. "Specific activation of Smad1 signaling pathways by the BMP7 Type 1 Receptor, ALK2," <i>J. Biol. Chem.</i> 273:25628-36. |
| | CE | Massague, et al. 2000. "Controlling TGF- β signaling," <i>Genes Deve.</i> 14:627-644. |
| | CF | Massague, J., 1998. "TGF- β signal transduction," <i>Ann. Rev. Biochem.</i> 67:753-791. |
| | CG | Miyazono, K., 2000. "TGF- β signaling by Smad proteins," <i>Cyto. Growth Factor Rev.</i> 11:15-22. |
| | CH | Murakami, et al. 1996. "Hypertensive and Hypotensive Mice produced by the introduction and disruption of genes on the Renin-Angiotensin system," <i>Blood Press. Suppl.</i> 2:36. |

| | |
|----|--|
| CI | Muralidhar, et al. 1993. "The Drosophila bendless gene encodes a neural protein related to ubiquitin-conjugating enzymes," Neuron 11:253-66. |
| CJ | Nakao, et al. 1997. "Identification of Smad7, a TGF β -inducible antagonist of TGF- β signalling," Nature 389:631-635. |
| CK | Nalefski, et al. 1996. "The C2 domain calcium-binding motif: Structural and functional diversity," Protein Sci. 5:2375-2390. |
| CL | Nefsky, et al. 1996. "Pul1 acts as an E6-AP-like protein ubiquitin ligase in the degradation of cdc25," Embo. J. 15:1301-1312. |
| CM | Patton, et al. 1998. "Combinatorial control in ubiquitin-dependent proteolysis: don't Skp the F-box hypothesis," Trends. Genet. 14:236-243. |
| CN | Plant, et al. 1997. "The c2 domain of the ubiquitin protein ligase Nedd4 mediates Ca ²⁺ -dependent plasma membrane localization," J. Biol. Chem. 272:32329-36. |
| CO | Pukatzki, et al. 1998. "A novel component involved in ubiquitination is required for development of Dictyostelium discoideum," J Biol. Chem. 273:24131-8. |
| CP | Reddi, A. H., 1998. "Role of morphogenetic proteins in skeletal tissue engineering and regeneration," Nature Biotech. 16:247-252. |
| CQ | Reeck, et al. 1987. "Homology" in proteins and nucleic acids: A terminology muddle and a way out of it," Cell 50:667. |
| CR | Rotin, D., 1998. "WW (WWP) domains: From structure to function," Curr. Topics Microbiol. Immunol. 228:115-133. |
| CS | Sasai, et al. 1997. "Ectodermal patterning in vertebrate embryos," Dev. Biol. 182:5-20. |
| CT | Scheiffner, et al. 1993. "The HPV-16 E6 and E6-AP Complex functions as a ubiquitin-protein ligase in the ubiquitination of p53," Cell 75:495-505. |
| CU | Staub, et al. 1997. "Immunolocalization of the ubiquitin-protein ligase Nedd4 in tissues expressing the epithelial Na ⁺ channel (EnaC)," Am. J Physiol. 272:C1871-80. |
| CV | Staub, et al. 2000. "Regulation of stability and function of the epithelial Na ⁺ channel (EnaC) by utiquitination," Kidney Int. 57:809-815 |
| CW | Staub, et al. 1997. "Regulation of the epithelial Na ⁺ channel by Nedd4 and ubiquitination," EMBO J 16:6325-6336. |
| CX | Staub, et al. 1996. "WW domains," Structure 4:495-499. |
| CY | Staub, et al. 1996. "WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na ⁺ channel deleted in Liddle's syndrome," EMBO J. 15:2371-2380. |
| CZ | Suzuki, et al. 1997. "Smad5 induces ventral fates in Xenopus embryo," Dev. Biol. 184:402-405. |
| DA | Takase, et al. 1998. "Induction of Smad6 mRNA by bone morphogenetic proteins," Biochem. Biophys. Res. Commun. 244:26-29. |
| DB | Thomsen, G.H., 1997. "Antagonism within and around the organizer: BMP inhibitors in vertebrate body patterning," Trends Genet. 13:209-211. |
| DC | Thomsen, G.H., 1996. "Xenopus mothers against decapentaplegic is an embryonic ventralizing agent that acts downstream of the BMP-2/4 receptor," Development 122:2359-66. |

| | | |
|--|----|--|
| | DD | Tsukazaki, et al. 1998. "SARA , a FYVE domain protein that recruits Smad2 to the TGF β receptor," Cell 95:799-791. |
| | DE | Ulloa, et al. 1999. "Inhibition of transforming growth factor- β /SMAD signalling by the interferon- γ /STAT pathway," Nature 397:710-713. |
| | DF | van Kerkhof, et al. 2000. "Endocytosis and degradation of the growth hormone receptor are proteasome-dependent," J. Biol. Chem. 275:1575-1580. |
| | DG | Wang, , et al. 1999. "Functional domains of the Rsp5 Ubiquitin-protein ligase," Mol. Cell Biol. 19:342-52. |
| | DH | Whitman, M., 1998. "Smads and early developmental signaling by the TGF β superfamily," Genes and Dev. 12:2445-2462. |
| | DI | Wigler, et al. 1979. "Transformation of mamalian cells with genes from procaryotes and eucaryotes," Cell 16:777-785. |
| | DJ | Wilson, et al. 1997. "Concentration-dependent patterning of the Xenopus ectoderm by BMP4 and its signal transducer Smad1," Dev. 124:3177-3184. |
| | DK | Wrana, et al. 2000. "Regulation of Smad activity," Cell 100:189-192. |
| | DL | Wrana, et al. 2000. "The Smad pathway," Cytokine & Growth Factor Reviews 11:5-13. |
| EXAMINER | | DATE CONSIDERED |
| Hope Robinson | | 9/30/04 |
| <p>*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.</p> | | |

RECEIVED

JUN 23 2003

TECH CENTER 1000/2000